

On the number of points in general position in the plane

József Balogh ^{*} József Solymosi [†]

April 19, 2017

Abstract

In this paper we study some Erdős type problems in discrete geometry. Our main result is that we show that there is a planar point set of n points such that no four are collinear but no matter how we choose a subset of size $n^{5/6+o(1)}$ it contains a collinear triple. Another application studies ε -nets in a point-line system in the plane.

We prove the existence of some geometric constructions with a new tool, the so-called Hypergraph Container Method.

1 Introduction

We say that a set of $n \geq d + 1$ points on the Euclidean space \mathbb{R}^d is in *general position* if no hyperplane contains $d + 1$ points. One might expect that if in a set no hyperplane contains $d + 2$ points then a large subset of the n points can be selected such that the points of this subset are in general position. Determining that in the worse case how many points in general position could be chosen - like many other similar problems - was initiated by Paul Erdős [11, 13]. We consider the planar ($d = 2$) case first. The best known upper and lower bounds are far from each other and even a sublinear upper bound was hard to achieve. Our primary goal is to prove the existence of a better construction for the upper bound. Furthermore, we apply similar methods to other problems as well.

ε -nets are important concepts in computational geometry and approximation theory in computer science. On a base set A of n elements and a family of subsets $\mathcal{F} \subset 2^A$ a set $E \subset A$ is an ε -net if for every $B \in \mathcal{F}$ if $|B| \geq \varepsilon n$ then $E \cap B \neq \emptyset$. It is expected that if \mathcal{F} has low complexity then one can find a small ε -net. In their seminal paper Haussler and Welzl [18] proved that for any range space with VC dimension d , there is an ε -net of size

^{*}Department of Mathematical Sciences, University of Illinois at Urbana-Champaign, Urbana, Illinois 61801, USA jobal@math.uiuc.edu. Research is partially supported by NSF DMS-1500121, NSA Grant H98230-15-1-0002, Arnold O. Beckman Research Award (UIUC Campus Research Board 15006) and TT 12 MX-1-2013-0006.

[†]Department of Mathematics, University of British Columbia 1984 Mathematics Road, Vancouver British Columbia V6T 1Z4, Canada solymosi@math.ubc.ca. Research was partially supported by NSERC, ERC Advanced Research Grant AdG. 321104 and by Hungarian National Research Grant NK 104183.

$O((d/\varepsilon) \log(d/\varepsilon))$. For the definitions and for further details we refer to [20], or Chapter 15 in [25], Chapter 10 in [22], or the survey paper of related problems [15]. Here we are concerned with a special case. Answering a question by Matoušek, Seidel, and Welzl [23], Alon [2] proved that the minimum possible size of an ε -net for point objects and line ranges in the plane is super-linear in $1/\varepsilon$. Alon used the density Hales-Jewett theorem for the construction, so his bound was just enough to break linearity. As he writes at the end of the paper “*The problem of deciding whether or not there are natural geometric range spaces of VC-dimension d in which the minimum possible size of an ε -net is $\Omega((d/\varepsilon) \log(1/\varepsilon))$ remains open. It seems plausible to conjecture that there are such examples, and even to speculate that this is the case for the range space of lines in the plane, for appropriately defined planar sets of points.*” We cannot quite answer this question however we are able to give a better bound, $(1/\varepsilon) \log^{1/3-o(1)}(1/\varepsilon)$, for the range space of lines. In a related work Pach and Tardos [26] constructed range spaces induced by axis-parallel rectangles in the plane, in which the size of the smallest ε -net is $\Omega(\frac{1}{\varepsilon} \log \log \frac{1}{\varepsilon})$. By a theorem of Aronov, Ezra, and Sharir [3], this bound is tight.

Here, we consider problems where points in planes are of interest. In a follow-up paper we consider high dimensional variants. Some variants of all of our methods work there as well, however there are more technical details, therefore it looked reasonable to handle them separately.

We will use the method of hypergraph containers for the proof. This useful method was recently introduced independently by Balogh, Morris and Samotij [6], and by Saxton and Thomason [32]. Roughly speaking, it says that if a hypergraph \mathcal{H} has a uniform edge distribution, then one can find a relatively small collection of sets, *containers*, covering all independent sets in \mathcal{H} . One can also require that the container sets span few edges only. In our applications this later condition guarantees that all container sets are small. The right geometric construction determines a hypergraph where all large subsets contain an edge (e.g. a collinear triple).

Note that the container method was applied in a similar way in the graph setting [24] and [31], and in additive combinatorics [4], and there are several more recent papers, here we just listed the ones which were motivating our work. The merit of our work is that we have realized that this machinery is useful in discrete geometry. Given this method, still it is possible that better results could be obtained, in case someone was able to find more useful hypergraphs, or prove a variant of a container theorem tailored to these particular hypergraphs.

We refer the readers to [4, 5, 6, 32] for more details and applications on the container method.

Notation. The $O(f(n))$, $w(g(n))$, $o(h(n))$ notation are standard, i.e., $k(n) = O(f(n))$ if there is a constant $C > 0$ that $k(n) \leq C \cdot f(n)$; $k(n) = w(g(n))$ if there is a constant $C > 0$ that $k(n) \geq C \cdot g(n)$; $k(n) = o(h(n))$ if for every constant $C > 0$ we have $k(n) \leq C \cdot h(n)$ for every n sufficiently large. We use logarithms of base 2, unless otherwise indicated. Throughout the paper we omit floors and ceilings whenever they are not crucial.

2 Results

2.1 Points in general position

For an n -element set, $S \subset \mathbb{R}^2$, let $S' \subset S$ be the largest subset of points in general position and let $\alpha(S) = |S'|$. We define

$$\alpha(n) := \min\{\alpha(S) : |S| = n \text{ and } S \text{ contains no four points in a line}\}.$$

Erdős [13] proposed the problem to determine $\alpha(n)$. Füredi [14] proved

$$\omega(\sqrt{n \log n}) \leq \alpha(n) = o(n). \quad (1)$$

The lower bound was obtained using a general result of Phelps and Rödl [28], who gave a lower bound on the independence number of partial Steiner Systems. Füredi [14] proved the upper bound using the Density Hales-Jewett Theorem (DHJ) of Katznelson and Furstenberg [16, 17]. In this application Füredi takes a random (generic) projection of the combinatorial cube of the combinatorial space $\{1, 2, 3\}^N$ to the plane. This point set has no collinear four-tuple, however for any $\delta > 0$ if N is large enough then any subset of points of size at least $\delta \cdot 3^N$ contains a collinear triple, a projection of a combinatorial line. The best known bound for DHJ follows from a recent proof by D.H.J. Polymath [29]. A subset of $\{1, 2, 3\}^N$ of density δ contains a combinatorial line if N is at least a tower of 2's of height $O(1/\delta^3)$. We improve the upper bound in (1) showing the existence of a point set containing no four points in a line such that its every subset of size $n^{5/6+o(1)}$ contains three points on a line.

Theorem 2.1. *As n goes to infinity,*

$$\alpha(n) \leq n^{5/6+o(1)}.$$

It is far from being clear, if $5/6$ is the right constant in the exponent, however it could be that the point set that we constructed is close to be best possible, at least it should be close to be optimal. Even then, it might have a better exponent, and only the limitation of our method is the reason that we cannot determine it.

2.2 ε -nets and weak ε -nets

Here, for a given small $\varepsilon^* > 0$, we shall construct a point set S in the plane, such that the smallest $R \subset S$, which covers every line containing at least $\varepsilon^*|S|$ points from $|S|$ has size at least $(1/\varepsilon^*) \log^{1/3-o(1)}(1/\varepsilon^*)$, i.e., we prove the next theorem.

Theorem 2.2. *Let ε^* be an arbitrary small constant. Then there exists a point system S in the plane, that if $T \subset S$ intersects each line containing at least $\varepsilon^*|S|$ points, i.e., T is an ε^* -net, then*

$$|T| \geq \frac{1}{2\varepsilon^*} \log^{1/3} \left(\frac{1}{\varepsilon^*} \right) \log \log \left(\frac{1}{\varepsilon^*} \right)^{-1}.$$

Note that the $O(\frac{1}{\varepsilon}(\log \frac{1}{\varepsilon}))$ upper bound holds for any abstract range space with fixed VC-dimension, see [18].

There is a variant of the ε -net problem where the hitting set T does not need to be a subset of the base set. Given a point set S in the plane, an T subset of the plane is a *weak- ε -net* if every line containing at least $\varepsilon|S|$ points of S contains a point of T . Note that T is not necessarily subset of S . We are interested proving the existence of an S , where the smallest point set which contains a point from every **line** which is incident to at least $\varepsilon|S|$ points of S is large. Alon [2] gave a lower bound to weak-nets as well, similar to the previous case using D.H.J.. We are also going to give a construction to bound the size of weak nets with an additional feature; it works in the projective plane as well.

Here we can prove only a similar, but weaker bound than in the non-weak case.

Theorem 2.3. *Let ε^0 be an arbitrary small constant. Then there is a $0 < \varepsilon^* < \varepsilon^0$ such that the following holds. There exists a point system S in the plane, that if T intersects each line containing at least $\varepsilon^*|S|$ points, i.e., T is a weak ε^* -net, then*

$$|T| \geq \frac{1}{10\varepsilon^*} \left(\log \log \frac{1}{\varepsilon^*} \right)^{1/10}.$$

2.3 Decomposing coverage of the plane

The point-line duality of the plane maps points into lines, lines into points that the incidences (and non-incidences as well) are preserved. Let \mathcal{F} be a family of sets over X . We say that \mathcal{F} is an *r -fold covering* of X , if for every $x \in X$ there are at least r sets in \mathcal{F} containing x . A family of sets \mathcal{F} over X is *cover-decomposable* if there exists an r such that any r -fold covering of X with members of \mathcal{F} can be decomposed into two coverings. Pach, Tardos and Tóth [27], using the Hales-Jewett Theorem, proved that when \mathcal{F} is the set of lines, and $X = \mathbb{R}^2$, then this family is not cover-decomposable.

In particular, for every r they proved the existence of a finite $X \subset \mathbb{R}^2$ and a finite family of lines \mathcal{F} , that every point of X was covered by at least r lines of \mathcal{F} , and \mathcal{F} was not decomposable into two coverings. Following their proof, one can read out a density version, i.e., an estimate on how large is the smallest covering subset of \mathcal{F} .

In Section 6 for every $c > 0$ and r we proved the existence of a point set S in the plane, with the property that any subset of S of size $c \cdot |S|$ contains a collinear r -tuple. Denote \mathcal{L} the collection of lines containing r points from S . The dual of S , say S^* , is a family of lines, the dual of \mathcal{L} , say \mathcal{L}^* , is a family of points, with the property that each point of \mathcal{L}^* is covered by exactly r lines of S^* . Additionally, for every subset of S^* of size at least $c|S^*|$ there is a point which is covered by r lines from S^* , i.e., it is uncovered by the complement of that subset.

With other words, we gave an alternative proof of the result of [27], with a better quantitative bound on c in function of size of S .

Mani-Lavistka and Pach [21] observed that the Lovász Local Lemma could be used for the following.

Theorem 2.4. *Assume that positive integers satisfy $e(T + 1) < 2^{r-1}$. Given a finite point system \mathcal{P} and a finite line system \mathcal{L} in the plane such that every point is covered at least r times, and every line is intersected (in a point from \mathcal{P}) by at most T lines. Then \mathcal{L} can be decomposed into two covering systems.*

Slight modification of the construction from the previous sections shows some limits of Theorem 2.4, unfortunately it is not clear what the proper order of magnitude of T should be.

Theorem 2.5. *Let r be a positive integer, and $T := r^{5r^2}$. Then there is finite point system \mathcal{P} and a finite line system \mathcal{L} in the plane that every point is covered at least r times, and every line is intersected (in a point from \mathcal{P}) by at most T lines from \mathcal{L} , and \mathcal{L} cannot be decomposed into two covering systems.*

For self-completeness we sketch the proof of Theorem 2.4: Form a hypergraph, with vertex set \mathcal{L} and edge set $\{F_x : x \in \mathcal{P}\}$, where F_x is the collection of lines containing x . By the Lovász Local Lemma it is 2-colorable if no F_x is intersected by more than T other hyperedges.

We prove the dual statement of Theorem 2.5. Note that here when we say that a collection of points \mathcal{P} cannot be decomposed into two covering systems of a family of lines \mathcal{L} , we mean that \mathcal{P} cannot be partitioned into \mathcal{P}_1 and \mathcal{P}_2 , such that every line of \mathcal{L} contains a point from both \mathcal{P}_1 and \mathcal{P}_2 .

Theorem 2.6. *Let r be a positive integer, and $T := r^{5r^2}$. There is finite line system \mathcal{L} and a finite point system \mathcal{P} in the plane that*

- (i) *every line contains at least r points,*
- (ii) *every point from \mathcal{P} shares a line from \mathcal{L} by at most T points from \mathcal{P} ,*
- (iii) *\mathcal{P} cannot be decomposed into two covering systems.*

2.4 Chromatic number of cell-arrangements in the plane

A simple arrangement of a set \mathcal{L} of lines in \mathbb{R}^2 is when there are no parallel lines and no three lines going through the same point. It decomposes the plane into the set \mathcal{C} of cells, i.e. maximal connected components of $\mathbb{R}^2 - \cup_{\ell \in \mathcal{L}} \ell$. In [9], the hypergraph $\mathcal{H}_{\text{line-cell}} = (\mathcal{L}; \mathcal{C})$ was defined, with the vertex set \mathcal{L} , and each hyperedge $F \in \mathcal{C}$ is formed by the set of lines forming the boundary of a cell of \mathcal{L} . It was proved in [9], that there is a family \mathcal{L} of n lines, that the corresponding hypergraph $\mathcal{H}_{\text{line-cell}}$ has chromatic number $\Omega(\log n / \log \log n)$, and in [1] that is always $O(\sqrt{n / \log n})$. In [1], a relation to the Erdős (3, 4)-problem was pointed out: Assume that given n points in a plane, no four in a line, and any t of them contains three on a line. Then one can take the dual of the point system to obtain a family of n lines, that no four of them intersect in a point, but any t of them contains three having a common point. A small perturbation of the lines changes every intersecting triplets of lines into a triangle, i.e., having a good construction for the Erdős (3, 4)-problem gives automatically a density version of [9]. To summarize, Theorem 2.1 instantly implies the following result.

Theorem 2.7. *There are n lines in the plane that any subset of it of size $n^{5/6+o(1)}$ completely contains a cell. In particular, the cell-chromatic number of the system is at least $n^{1/6-o(1)}$.*

3 The hypergraph container theorem

First, we present our main tool, the hypergraph container theorem. Let \mathcal{H} be an r -uniform hypergraph with average degree d . For every $S \subseteq V(\mathcal{H})$, its co-degree, denoted by $d(S)$, is the number of edges in \mathcal{H} containing S , i.e.,

$$d(S) = |\{e \in E(\mathcal{H}) : S \subseteq e\}|.$$

For every $j \in [r]$, denote by Δ_j the j -th maximum co-degree, i.e.,

$$\Delta_j = \max\{d(S) : S \subseteq V(\mathcal{H}), |S| = j\}.$$

For $\tau \in (0, 1)$, define

$$\Delta(\mathcal{H}, \tau) = 2^{\binom{r}{2}-1} \sum_{j=2}^r \frac{\Delta_j}{d\tau^{j-1}2^{\binom{j-1}{2}}}.$$

In particular, when $r = 3$

$$\Delta(\mathcal{H}, \tau) = \frac{4\Delta_2}{d\tau} + \frac{2\Delta_3}{d\tau^2}.$$

We are going to use the following version of the hypergraph container theorem (Corollary 3.6 in [32]).

Theorem 3.1. *Let \mathcal{H} be an r -uniform hypergraph on vertex set $[N]$. Let $0 < \varepsilon, \tau < 1/2$. Suppose that $\tau < 1/(200 \cdot r \cdot r!^2)$ and $\Delta(\mathcal{H}, \tau) \leq \varepsilon/(12r!)$. Then there exists $c = c(r) \leq 1000 \cdot r \cdot r!^3$ and a collection of vertex subsets \mathcal{C} such that*

- (i) *every independent set in \mathcal{H} is a subset of some $A \in \mathcal{C}$;*
- (ii) *for every $A \in \mathcal{C}$, $e(\mathcal{H}[A]) \leq \varepsilon \cdot e(\mathcal{H})$;*
- (iii) *$\log |\mathcal{C}| \leq cN\tau \cdot \log(1/\varepsilon) \cdot \log(1/\tau)$.*

4 Supersaturation of collinear point sets in the grid.

Given integers $n, k, r \geq 3$, we consider the following r -uniform hypergraph $\mathcal{H} = \mathcal{H}(n, k, r)$ encoding the set of all collinear r -tuples in $[n]^k = \{1, 2, \dots, n\}^k$, we let $V(\mathcal{H}) = [n]^k$ and the edge set of \mathcal{H} consists of all collinear r -tuples.

While it is easy to compute the number of vertices of \mathcal{H} , $|V(\mathcal{H})| = n^k =: N$, it takes more effort to estimate the number of edges. For the bounds below, we shall assume that $r, k \leq 0.01 \cdot \log n$.

Claim 4.1. (i) If $r \leq k$ then

$$\frac{n^{2k}}{r^{2k}} \leq e(\mathcal{H}) \leq \frac{k \cdot 2^{r+k}}{r!} n^{2k}. \quad (2)$$

(ii) If $r = k + 1$ then

$$\frac{n^{2k}}{r^{2k}} \leq e(\mathcal{H}) \leq \frac{k \cdot 2^{r+k}}{r!} \cdot n^{2k} \log n. \quad (3)$$

(iii) If $k + 1 < r \leq 2k$ then

$$k \cdot \binom{n}{r} \cdot n^{k-1} \leq e(\mathcal{H}) \leq \frac{k \cdot 2^{r+k+1}}{r!} \cdot n^{r+k-1}. \quad (4)$$

Proof: For the lower bounds, we show the existence of many collinear r -tuples. A collinear r -tuple can be written in a form $u, u + v, \dots, u + (r - 1)v$ for some $u, v \in \mathbb{Z}^k$. In order to keep the points in $[n]^d$, we shall choose $u, v \in [n/r]^k$, which proves (i) and (ii). For (iii), when $r > k$, we can do better, as in any of the k (axis-parallel) directions we have n^{k-1} parallel lines, each containing n points, i.e. $\binom{n}{r}$ collinear r -tuples.

For the upper bound we need more care. Write $L(t)$ for the number of lines containing more than 2^t and at most 2^{t+1} points. Then we want to bound

$$\sum_{t=1}^{\log n} L(t) \cdot \binom{2^{t+1}}{r}. \quad (5)$$

If a line contains exactly ℓ points, where $2^t < \ell \leq 2^{t+1}$, then we can list these ℓ points as $u, u + v, \dots, u + (\ell - 1)v$ for some $u, v \in \mathbb{Z}^k$. Observe that because these points are in $[n]^k$, the absolute value of each coordinate of v is at most $n/(\ell - 1)$. By symmetry, and for the price of a factor of 2^k , we can assume that each coordinate of v is non-negative. This implies that at least one coordinate of u is at most $n/(\ell - 1)$, otherwise $u - v$ was on the line as well. Using that $2^t \leq \ell < 2^{t+1}$ this means that at least one coordinate of u is at most $n/2^t$, and each coordinate of v is at most $n/2^t$, meaning that the number of choices for v is at most $(n/2^t)^k$, and for u is at most $k \cdot n^k/2^t$, therefore $L(t) \leq 2^k \cdot (n/2^t)^k \cdot k \cdot n^k/2^t$. Writing this into (5) we obtain the following upper bound on the number of collinear r -tuples:

$$\sum_{t=1}^{\log n} 2^k \cdot (n/2^t)^k \cdot k \cdot n^k/2^t \cdot \binom{2^{t+1}}{r} \leq \sum_{t=1}^{\log n} \frac{k \cdot 2^{r+k}}{r!} n^{2k} 2^{t(r-k-1)}. \quad (6)$$

When $r \leq k$ then $\sum_t 2^{t(r-k-1)} \leq 1$. When $r = k + 1$, then $\sum_t 2^{t(r-k-1)} \leq \log n$. If $r > k + 1$ then $\sum_t 2^{t(r-k-1)} \leq 2 \cdot n^{r-k-1}$. Q.E.D.

A key ingredient is the supersaturation lemma, first we consider the $k = r = 3$ case.

Lemma 4.2. In $\mathcal{H}(n, 3, 3)$ every set of vertices of size n^{3-s} spans at least $\frac{n^{6-4s}}{3 \cdot 10^9 \log n}$ hyperedges, where $0 \leq s < 1$ and n is sufficiently large.

Proof. Let S be an arbitrary subset of vertices of size $|S| = n^{3-s}$, where $0 \leq s < 1$. Let $t = 2000n^s$ and

$$U := \{(a, b, c) : 1 \leq a \leq n/t, \quad -n \leq b, c \leq n\},$$

$$V = \{(a, b, c) : a \geq |b|, |c|, \quad n/t \leq a \leq 2n/t, \quad a \text{ is a prime}\}.$$

We define the collection of lines $\mathcal{L} = \mathcal{L}(t)$, whose starting points are in U , and their directions are in V (note that we define these lines in \mathbb{R}^3 , and we are interested in their intersection with $[n]^3$). The line system has several properties.

Claim 4.3. (i) Every line in \mathcal{L} contains at most t points from $[n]^3$.

(ii) $|\mathcal{L}| \leq 300n^6/(t^4 \cdot \log(n/t))$.

(iii) Every point in $[n]^3$ is contained in at least $n^3/(t^3 \log(n/t))$ and at most $50n^3/(t^3 \cdot \log(n/t))$ lines from \mathcal{L} .

Proof. Part (i) follows, as the absolute value of the first coordinate of a direction is at least n/t . Part (ii) follows from considering all choices from U and V , and that the number of prime numbers smaller than x is at least $0.9x/\log x$ and at most $1.1x/\log x$ for x sufficiently large. For (iii), the lower and upper bounds are coming from the size of V . Q.E.D.

The number of point-line incidences of the points from S and lines from \mathcal{L} by Claim 4.3 (iii) is at least $n^{3-s} \cdot n^3/(t^3 \log(n/t))$, hence the average number of points per line is at least $t \cdot n^{-s}/300 > 6$. Therefore, the number of collinear triplets in S is at least

$$\binom{t \cdot n^{-s}/300}{3} \cdot \frac{300n^6}{t^4 \cdot \log(n/t)} \geq \frac{n^{6-3s}}{1100000t \log(n/t)} \geq \frac{n^{6-4s}}{3 \cdot 10^9 \log(n/t)}.$$

Q.E.D.

In the next supersaturation result on $\mathcal{H} = \mathcal{H}(n, k, r)$, we have $r = k$, and we are interested to find many edges in large subsets of vertices. The idea of the proof is similar to the one in Claim 4.3, the computation is somewhat more technical. Unfortunately, the result is useful only when $k < 0.01 \log^{1/2} n$, because of the 2^{3k^2} term, it might be useful to extend the proof to improve the exponent $1/2$ to $1 - c$ for some small constant $c > 0$.

Lemma 4.4. Let $0.1 < \gamma \leq 1$ and $r = k < 0.01 \log^{1/2} n$ and n be sufficiently large. Then every set of vertices of size γn^k spans at least

$$\frac{\gamma^{k+1} \cdot n^{2k}}{k! \cdot 2^{3k^2} \cdot 10 \cdot 2^{3k} \cdot k \cdot \log n}$$

hyperedges.

Proof. Let S be an arbitrary subset of vertices of size $|S| = \gamma n^k$ and fix $t = 10 \cdot 2^{2k} \cdot k/\gamma$. Let

$$U := \{(a_1, \dots, a_k) : 1 \leq a_1 \leq n/t, \quad -n \leq a_2, \dots, a_k \leq n\},$$

$$V = \{(a_1, \dots, a_k) : |a_1| \geq |a_2|, \dots, |a_k|, \quad n/t \leq a_1 \leq 2n/t, \quad a_1 \text{ is a prime}\}.$$

We define a collection of lines $\mathcal{L} = \mathcal{L}(t)$, whose starting points are in U , and their directions are in V . The line system has several properties.

Claim 4.5. (i) Every line in \mathcal{L} contains at most t points from $[n]^k$.

(ii) $|\mathcal{L}| \leq 2^{3k} n^{2k} / (t^{k+1} \cdot \log(n/t))$.

(iii) The number of lines in \mathcal{L} containing a given point is at least $2^k \cdot n^k / (t^k \log(n/t))$ and at most $4^{k+1} n^k / (t^k \cdot \log(n/t))$.

Proof. Part (i) follows, as the absolute value of the first coordinate of a direction is at least n/t . Part (ii) follows from considering all choices from U and V , i.e.,

$$|\mathcal{L}| \leq |U| \cdot |V| \leq (2n+1)^{k-1} (n/t) (4n/t+1)^{k-1} (3n/t) / \log(n/t) \leq \frac{2^{3k} \cdot n^{2k}}{t^{k+1} \cdot \log(n/t)},$$

by the Prime Number Theorem for n sufficiently large. For (iii), the lower and upper bounds are coming from the size of V . Q.E.D.

The number of point-line incidences of the points from S and lines from \mathcal{L} by Claim 4.5 (iii) is at least $\gamma \cdot 2^k \cdot n^{2k} / (t^k \log(n/t))$, hence the average number of points per line is at least $\gamma \cdot 2^k \cdot n^{2k} / (|\mathcal{L}| \cdot t^k \log(n/t)) \geq 2^{-2k} \cdot \gamma \cdot t$. Therefore, using that $t > 10 \cdot 2^{2k} \cdot k / \gamma$, the number of collinear r -tuples in S is at least

$$\binom{\gamma \cdot 2^k \cdot n^{2k} / (|\mathcal{L}| \cdot t^k \log(n/t))}{r} \cdot |\mathcal{L}| \geq \binom{2^{-2k} t \cdot \gamma}{k} \cdot \frac{2^{3k} n^{2k}}{t^{k+1} \cdot \log(n/t)} \geq \frac{\gamma^k \cdot n^{2k}}{k! \cdot 2^{3k^2} \cdot t \cdot \log n}.$$

Q.E.D.

5 The container method on $[n]^3$: Proof of Theorem 2.1

We apply the container lemma several times to the hypergraph $\mathcal{H} = \mathcal{H}(n, 3, 3)$ and to its subhypergraphs. First we let $C^0 := [n]^3$, $\mathcal{C}^0 := \{C^0\}$ and write $\mathcal{H}^0 = \mathcal{H}[C^0] = \mathcal{H}$. Applying the container lemma to \mathcal{H}^0 will output a container family \mathcal{C}^1 . For each $C_j^1 \in \mathcal{C}^1$ we consider the hypergraph $\mathcal{H}_j^1 := \mathcal{H}[C_j^1]$ and we apply the container lemma. The collection of all container sets obtained from all $C_j^1 \in \mathcal{C}^1$ will form \mathcal{C}^2 . We iterate this process, i.e. we apply the container lemma for the hypergraphs spanned by the containers and will continue it, until each of the container sets will be sufficiently small. We will prove that the number of steps is $O(1)$, therefore the total number of containers will be still small. Note that one could form a general container theorem for hypergraphs with a strong supersaturation property along these ideas, but somehow it looks like that it is better to work on the particular hypergraphs separately. Similar multiphase applications of the container method appeared for example in [7] and [10].

In the final step of the proof of the theorem, as usual in the container method, using the union bound, we could claim that none the container sets contains too many elements of a random subset of the vertices.

We fix an arbitrary small constant $f > 0$, this will show up in the error term in the exponent next to $5/6$, also the number of iterations will be $O(1/f)$. For the generic iteration step i , assume that we work with $\mathcal{H}_j^i \subset \mathcal{H}$ spanned by a container set $C_j^i = V(\mathcal{H}_j^i)$, where i stands for the number of rounds and j is just a running label over various hypergraphs. Define $0 \leq s = s(i, j)$ such that

$$v(\mathcal{H}_j^i) = n^{3-s}.$$

Note, that in the first step we have $s = 0$. If $s \geq 1/3 - f$ then C_j^i is sufficiently small, and we put it into the final container family \mathcal{C} , and we shall not do anything with it.

Now we apply the container lemma on $\mathcal{H}_j^i = \mathcal{H}[C_j^i]$: By Lemma 4.2 we have a lower bound on $e(\mathcal{H}_j^i)$, hence on the average degree of \mathcal{H}_j^i as well:

$$e(\mathcal{H}_j^i) \geq \frac{n^{6-4s}}{3 \cdot 10^9 \log n} \quad \text{and} \quad d(\mathcal{H}_j^i) \geq \frac{3n^{6-4s}}{3 \cdot 10^9 \log n \cdot n^{3-s}} = \frac{n^{3-3s}}{10^9 \log n}.$$

Observe that the hypergraph \mathcal{H}^0 , and hence each subhypergraph of it satisfies that

$$\Delta_2 < n, \quad \Delta_3 = 1.$$

We set

$$\tau = n^{s-4/3} \quad \text{and} \quad \varepsilon = n^{s-1/3+f/2} < n^{-f/2}.$$

To apply Theorem 3.1 we need the following condition to be satisfied (when n is sufficiently large)

$$\Delta(\mathcal{H}_j^i, \tau) = \frac{4\Delta_2}{d\tau} + \frac{2\Delta_3}{d\tau^2} \leq \frac{4 \cdot 10^9 \cdot \log n}{n^{2/3-2s}} + \frac{2 \cdot 10^9 \cdot \log n}{n^{1/3-s}} \leq \frac{\varepsilon}{72}.$$

By Theorem 3.1, there is a family \mathcal{C}_j^{i+1} of containers such that

$$|\mathcal{C}_j^{i+1}| \leq 2^{10^6 \cdot n^{3-s} \tau \log^2 n} = 2^{10^6 \cdot n^{5/3} \log^2 n},$$

and each $C \in \mathcal{C}_j^{i+1}$ spans at most $\varepsilon \cdot e(\mathcal{H}_j^i)$ edges.

In the next round, for each $C \in \mathcal{C}_j^{i+1}$ we run the same process (say C is playing the role of $V(\mathcal{H}_{j'}^{i+1})$). The key point of the iteration is that at each iteration step, when i is increasing to $i+1$, then either $e(\mathcal{H}_j^i) \leq n^{14/3+f}$ and $v(\mathcal{H}_j^i) \leq n^{8/3+f}$ or the number of edges is shrinking with a factor of at least $n^{-f/2}$, therefore the number of iteration steps is at most $12/f$, hence the total number of containers will be at most

$$2^{(12/f)10^7 \cdot n^{5/3} \log^2 n}.$$

Here, the key step was that when we are applying the container lemma to a hypergraph spanned by a container, the number of new containers is not increasing a lot, just by a constant factor in the exponent, which accumulates only to a constant factor in the exponent, as the number of iterations was a constant. The iteration stops, i.e., we do not touch a

container C when it is small, i.e., $|C| \leq n^{8/3+f}$.

To summarize: we obtained a family of containers \mathcal{C} containing at most $2^{(12/f)10^7 \cdot n^{5/3} \log^2 n}$ sets, each of size at most $n^{8/3+f}$.

Now we turn to the probabilistic construction of our point set in the plane. We take a p -random subset of $[n]^3$, keeping with probability p a point, independently of the other points. Having this random point set, we remove a point from each collinear 4-tuple. We want to choose a maximum value of p that most of the randomly chosen points should stay. The expected number of random points is pn^3 , by Claim 4.1 (ii) the expected number of collinear 4-tuples is at most $100 \cdot p^4 \log n \cdot n^6$, hence for

$$p = \frac{1}{n \log^{1/2} n}$$

we have

$$pn^3 \gg 100p^4 \log n \cdot n^6.$$

Denote the resulted set by S .

By the container method above we proved that there are at most $2^{(12/f)10^7 \cdot n^{5/3} \log^2 n}$ containers (subsets of the vertex set), each set having size at most $n^{8/3+f}$, and each independent set is contained in one of them.

We claim, using the first moment method, that S unlikely will contain an independent set of size m when $m = n^{5/3+f}$:

$$2^{(12/f)10^7 \cdot n^{5/3} \log^2 n} \cdot \binom{n^{8/3+f}}{m} p^m \leq 2^{(12/f)10^7 \cdot n^{5/3} \log^2 n} \cdot \left(\frac{e}{\log^{1/2} n} \right)^m = o(1).$$

Therefore, there is a set S which does not contain an independent set of size $n^{5/3+f}$, does not contain 4 points in a line, and

$$\frac{n^2}{2 \log^{1/2} n} \leq |S| \leq \frac{2n^2}{\log^{1/2} n}.$$

The set S can be projected into the plane that collinear point tuples stay on a line, and no new collinear point tuples are created. Thus we have about $n^2 / \log^{1/2} n$ points in the plane that every subset of size $n^{5/3+f}$ contains three on a line. As $f > 0$ was an arbitrary small constant this completes the proof of Theorem 2.1. Q.E.D.

6 Container method on $[n]^r$: applying to ε -nets

In the beginning of the proof we apply the container method as in the previous section, just to a different hypergraph. We complete the proof using ideas of Alon [2]. Note that here the application of the container method is somewhat simpler (though the supersaturation part is more technical), because it is sufficient to obtain containers of size $0.1 \cdot n^r$, i.e., 1/10th

proportion of the vertex set of the hypergraph, therefore it is sufficient to apply the container lemma only once.

Here, we shall work with $\mathcal{H} = \mathcal{H}(n, r, r)$, where $r \leq 0.01 \log^{1/2} n / \log \log n$, but $r = r(n) \gg \log \log n$, we assume that both r and n are sufficiently large, that the computations hold in this section, in particular Lemma 4.4 holds. Using that there are at most n points in a line, we have

$$\Delta_2 = \binom{n-2}{r-2}, \quad \dots, \quad \Delta_i = \binom{n-i}{r-i}, \quad \dots, \quad \Delta_r = 1.$$

Using Claim 4.1 (i) we can estimate the average degree of \mathcal{H}

$$\frac{n^{2r}}{r^{2r}} \leq e(\mathcal{H}) \leq \frac{2^{2r}}{(r-1)!} n^{2r} \quad \text{and} \quad \frac{n^r}{r^{2r-1}} \leq d = d(\mathcal{H}) \leq \frac{r^2 \cdot 2^{2r}}{r!} n^r.$$

Also, by Claim 4.1 (ii), the number of collinear $(r+1)$ -tuples is at most $\frac{(r+1) \cdot 2^{2r+1}}{r!} n^{2r} \cdot \log n$.

We need the following supersaturation statement, which is implied by Lemma 4.4.

Lemma 6.1. *Let $T \subset V(\mathcal{H})$ with $|T| \geq 0.1v(\mathcal{H})$ then*

$$e(\mathcal{H}[T]) \geq \frac{n^{2r}}{10^{r+2} 2^{3r^2+3r} (r+1)! \cdot \log n} \geq 10^{-r^2} e(\mathcal{H}).$$

Now we take a random subset S of $v(\mathcal{H})$, choosing each vertex with probability p independently of the other choices. We want that the number of collinear $(r+1)$ -tuples should be less than the number of vertices, so removing one point from each collinear $(r+1)$ -tuple results in a set with containing no collinear $(r+1)$ -tuple, so we need

$$pn^r \gg p^{r+1} \frac{(r+1) \cdot 2^{2r+1}}{r!} n^{2r} \log n,$$

observing that $\log^{1/r} n = 1 + o(1)$, it is satisfied when

$$p = \frac{r}{20n}. \tag{7}$$

We set $\varepsilon = 10^{-r^2}$, which by Lemma 6.1 means that if a container spans less than $\varepsilon \cdot e(\mathcal{H})$ edges, then it has at most $n^r/10$ vertices.

For the application of the container lemma, we need to choose τ satisfying the conditions below:

$$\tau < \frac{1}{200 \cdot r \cdot r!^2} \quad \text{and} \tag{8}$$

$$\Delta(\mathcal{H}, \tau) = 2^{\binom{r}{2}-1} \sum_{j=2}^r \frac{\Delta_j}{d\tau^{j-1} 2^{\binom{j-1}{2}}} \leq 2^{\binom{r}{2}-1} \sum_{j=2}^r \frac{\binom{n-j}{r-j} \cdot r^{2r-1}}{n^r \tau^{j-1} 2^{\binom{j-1}{2}}} \leq \frac{\varepsilon}{12r!}. \tag{9}$$

To satisfy the above conditions we choose

$$\tau = n^{-1-0.9/(r-1)}.$$

Maybe, it is useful to give a heads up, why it is a good choice. The inequality (8) is easily satisfied as $r \leq \log^{1/2} n$. The bulk of (9) is that $n^{-r} \tau^{1-j} \ll 10^{-r^2}$, which is the hardest to satisfy when $r = j$. Plugging our choice of τ simplifies it to $n^{0.1} \gg 10^{r^2}$, which is satisfied as $r \ll \log^{1/2} n$.

The Container Theorem, Theorem 3.1 gives the existence of at most

$$2^{1000 \cdot r \cdot r!^3 v(\mathcal{H}) \tau \cdot \log(1/\varepsilon) \cdot \log(1/\tau)} \leq 2^{n^{-1-0.8/r} \cdot v(\mathcal{H})}$$

containers, each of size at most

$$0.1 \cdot v(\mathcal{H}),$$

that each independent set is contained in one of them. Note that the above inequality is satisfied as $n^{0.1/r} \gg r^r \cdot \log n$ for our choice of r , actually this is from where the upper bound for r is coming from, though we used at many other places that $r \leq \log^{1/2} n$.

Now we can give an upper bound on the number of independent sets in the p -random subset of \mathcal{H} of size $m = 0.45pv(\mathcal{H})$, simply multiplying the number of containers with the number of choices of an m -set from a container:

$$2^{n^{-1-0.8/r} \cdot v(\mathcal{H})} \cdot \binom{0.1 \cdot v(\mathcal{H})}{m} \cdot p^m \leq 2^{n^{-1-0.8/r} \cdot v(\mathcal{H})} (0.61)^m < 2^{(n^{-1-0.8/r} - p/5) \cdot v(\mathcal{H})} = o(1).$$

Now we can put together the proof: Consider a p -random set of $v(\mathcal{H})$. By the choice of p , the removal of $o(pv(\mathcal{H}))$ points can make it to have the property that no $r+1$ points are in a line. Project the set into the plane, keeping the collinearity of the points, (more importantly not creating new collinear point sets) and denote the resulted set by S . Note that we had that with high probability, S does not contain a subset of size $|S|/2$ which contains no r points in a line. Now we choose carefully the parameters: Let ε^* be a small positive constant, and set

$$r := \log^{1/3} \left(\frac{1}{\varepsilon^*} \right) \log \log \left(\frac{1}{\varepsilon^*} \right)^{-1} \quad \text{and} \quad |S| = \frac{r}{\varepsilon^*}.$$

One can compute n , using (7) and $|S| = p \cdot n^r$, but we do not need explicit formula for n , existence of it is sufficient for us for every small $\varepsilon^* > 0$, maybe useful to state that

$$\varepsilon^* = 2^{-(1+o(1))r^3}, \quad n = 2^{(1+o(1))r^2}, \quad p = 2^{-(1+o(1))r^2}.$$

If $T \subset S$ is an ε^* -net, then it has to intersect every line containing r points, which means that $S - T$ is an independent set. Note that this is the part of the proof where it is used that there is no line containing $r+1$ points. There is no independent set of size $|S|/2$ in S , so $|S| - |T| < |S|/2$, which means that

$$|T| \geq \frac{|S|}{2} \geq \frac{1}{2\varepsilon^*} \log^{1/3} \left(\frac{1}{\varepsilon^*} \right) \log \log \left(\frac{1}{\varepsilon^*} \right)^{-1}.$$

Q.E.D.

7 Weak ε -nets

We are also going to give a construction to bound the size of weak nets with an additional feature; it works in the projective plane as well. The projection of $[k]^n$ has a very small weak ε -net if one can use points in infinity, i.e., if we consider the problem in the projective plane. There are $2^n - 1$ different directions for the lines, so $2^n - 1$ points can cover all k -rich lines. In our construction after the tilting and projection the rich lines are in general position.

Here we can prove only a similar, but weaker bound than in the non-weak case. Our main trouble is that the points are in too many lines, and there are far too many eliminated points to handle. Still we have a similar approach, but we consider a much sparser hypergraph. Denote $\mathcal{F} = \mathcal{F}(n, k, r)$ the following hypergraph. Its vertex set is $[k]^n$, and an r -tuple of vertices forms a hyperedge if they are on an axis-parallel line, i.e., all but one coordinate of their points are the same.

As before, we shall project it to the plane. We are using a map, which keeps 3 or more points collinear, if they were on an axis-parallel line, and no new collinear 3-tuples are created. Here we have a technical attempt to describe this map, which could be skipped by the reader. To avoid non axis-parallel triples to be collinear we use a generic Cartesian product. Let $H_i = \{a_1, a_2, \dots, a_n\}$ where all n^2 entries, a_j -s, are algebraically independent, distinct, non-zero real numbers. Now let $V(\mathcal{H})$ denote the Cartesian product $H_1 \times H_2 \times \dots \times H_n$. In this "grid" only axis-parallel triples are collinear. For the weak ε -nets application in the projective plane this construction is not applicable yet. The problem is that all axis-parallel lines are incident to only a few, n , points in the infinity. The construction below is described in the real space, \mathbb{R}^n , but it has a natural embedding into the projective space, $\mathbb{P}\mathbb{R}^n$. We will apply two maps to avoid lines being parallel. First we show the construction in \mathbb{R}^3 . This "un-parallelization" technique was used by Kollár in [19].

Let's define a map, $\nu_3 : \mathbb{R}^3 \rightarrow \mathbb{R}^7$, such that

$$\nu_3 : (x_1, x_2, x_3) \mapsto (x_1, x_2, x_3, x_1 \cdot x_2, x_1 \cdot x_3, x_2 \cdot x_3, x_1 \cdot x_2 \cdot x_3).$$

A parametric equation of a line which is parallel to the x_1 axis, say, is $[t+a, b, c]$, and its image is $[a, b, c, ba, ca, cd, bca] + t[1, 0, 0, b, c, 0, bc]$. As the map is injective, non-crossing lines remain non-crossing. Let's check two parallel lines, ℓ_i and ℓ_j . After the map the two direction vectors are $[1, 0, 0, b_i, c_i, 0, b_i c_i]$ and $[1, 0, 0, b_j, c_j, 0, b_j c_j]$. If the vectors are parallel then there is a non-zero multiplier $\alpha \in \mathbb{R}$, such that $[b_i, c_i, b_i c_i] = \alpha[b_j, c_j, b_j c_j]$. Then $b_i c_j = b_j c_i$ which would contradict to the algebraic independence of the elements of H_i .

For the general case let us apply a map, $\nu_n : \mathbb{R}^n \rightarrow \mathbb{R}^{2^n-1}$, where a point is mapped to all possible products of its coordinates.

$$\nu : (x_1, x_2, \dots, x_n) \mapsto (x_1, x_2, \dots, x_n, x_1 \cdot x_2, \dots, x_{n-1} \cdot x_n, \dots, x_1 \cdot x_2 \cdot \dots \cdot x_n).$$

Only the axis parallel collinear triples will stay collinear in the image of $V(\mathcal{H})$ in \mathbb{R}^{2^n-1} under the map ν_d . Now we can project the image back generically to \mathbb{R}^n . Other maps, like Veronese embeddings, would be sufficient too.

The important parameters of \mathcal{F} are the following:

$$v(\mathcal{F}) = k^n, \quad e(\mathcal{F}) = n \binom{k}{r} k^{n-1}, \quad d(\mathcal{F}) = \frac{rn}{k} \binom{k}{r}, \quad \Delta_i = \binom{k-i}{r-i}.$$

In the proof below we chose the parameters

$$n = k^r, \quad r = (\log n)^{1/5}, \quad r = t^2.$$

These choices are not optimal, in particular the exponent $1/5$, we aimed for clarity rather than pushing the proof to the best possible result. We emphasize that in this section if we say that a set of points of $[k]^n$ is collinear, then they are in an axis-parallel line of $[k]^n$. We need the following supersaturation lemma:

Lemma 7.1. *Let $\gamma > 10r/k$ and $T \subset V(\mathcal{F})$ with $|T| > \gamma v(\mathcal{F})$. Then*

$$e(\mathcal{F}[T]) \geq (\gamma - r/k)^r e(\mathcal{F}).$$

Proof. Fix a coordinate (can be done n ways), a direction (or a representative) of a line given by the other coordinates (k^{n-1} ways). In average, a line contains at least γk points from T , having $\binom{\gamma k}{r}$ collinear r tuples. Using the convexity of the binomial coefficients, the number of hyperedges in T is at least

$$n \cdot k^{n-1} \cdot \binom{\gamma k}{r} = \frac{\binom{\gamma k}{r}}{\binom{k}{r}} e(\mathcal{F}) \geq \left(\frac{\gamma k - r}{k} \right)^r e(\mathcal{F}).$$

Q.E.D.

The construction of the set of points is as before; first we choose a random p -subset of the grid $[k]^n$, then we remove a point from each collinear $(r+1)$ -tuples. We shall do an additional sparsening, for every collection of t lines, each containing r (p -random) points, and intersecting in one not p -random point, we remove one out of the tr points. With other words, after this sparsening there will be no t collinear (p -random) r -tuples which could be covered by one non- p -random point. We need to choose p carefully, as we do not want to remove too many points, these restrictions will give an upper bound on p .

Using similar method as in the ε -net proof, we shall prove that the obtained set will have no large independent set. From here, we shall obtain a lower bound on p . Using a switching idea of Alon [2], we will conclude that no small weak ε -net exists.

The first condition on p is coming from that the expected number of collinear $(r+1)$ -tuples needs to be much less than the expected number of points, i.e., $p^{r+1} n \binom{k}{r+1} k^{n-1} \ll p \cdot k^n$, which is satisfied when

$$p \ll \frac{r}{k \cdot n^{1/r}} = \frac{r}{k^2}. \quad (10)$$

For the second condition we use a key property of the projection (that Alon [2] used as well) that if there are at least 3 collinear r -tuples of points passing through the same point, then this point was a grid point(!). This aids us in the counting of the t -tuples of lines -each

containing r random grid points, as we can start by their intersection point. The expected number of t -tuples of such lines is, using that each point is in n lines

$$k^n \cdot \binom{n}{t} \binom{k}{r}^t p^{rt},$$

where k^n is the number of points, $\binom{n}{t}$ is the number of ways to choose the t directions, $\binom{k}{r}^t$ the number of ways to choose on each r points, that all were kept with probability p each.

We need that it is much less than $p \cdot k^n$, which is satisfied if

$$p \ll \frac{1}{k^{1+1/(rt-1)} \cdot n^{t/(rt-1)}}. \quad (11)$$

For the container lemma we set

$$\tau = \frac{r \cdot 2^r}{k \cdot n^{1/(r-1)}}, \quad \varepsilon = 10^{-(r+1)}.$$

In order to apply Theorem 3.1, we have to check the condition on $\Delta(\mathcal{F}, \tau), \tau, \varepsilon$:

$$\Delta(\mathcal{F}, \tau) = 2^{\binom{r}{2}-1} \sum_{j=2}^r \frac{\Delta_j}{d\tau^{j-1} 2^{\binom{j-1}{2}}} \leq 2^{\binom{r}{2}-1} \sum_{j=2}^r \frac{\binom{k-j}{r-j} \cdot k}{r \cdot n \cdot \binom{k}{r} \cdot \tau^{j-1} 2^{\binom{j-1}{2}}} \leq \frac{\varepsilon}{12r!},$$

which is satisfied by our choice of parameters. Note that when $j > r/2$ then $2^{\binom{j}{2}+r(j-1)}$, and if $j \leq r/2$ then $n^{(r-j)/(r-1)}$ is the dominant term in the denominator, which makes the relation true.

The Container Theorem 3.1 gives the existence of at most

$$2^{1000 \cdot r \cdot r!^3 v(\mathcal{F}) \tau \cdot \log(1/\varepsilon) \cdot \log(1/\tau)} \leq 2^{v(\mathcal{F}) r^{4r} \cdot k^{-1} \cdot n^{-1/(r-1)}}$$

containers, that each independent set is contained in one of them, that each spans at most $\varepsilon \cdot e(\mathcal{F})$ hyperedges. By Lemma 7.1, each container has at most $v(\mathcal{F})/9$ vertices.

Now we can give an upper bound on the number of independent sets in the p -random subset of \mathcal{H} of size $m = 2pv(\mathcal{H})/3$, simply multiplying the number of containers with the number of choices of an m -set from a container:

$$2^{v(\mathcal{F}) r^{4r} \cdot k^{-1} \cdot n^{-1/(r-1)}} \cdot \binom{v(\mathcal{H})/9}{m} \cdot p^m \leq 2^{v(\mathcal{F}) r^{4r} \cdot k^{-1} \cdot n^{-1/(r-1)}} 2^{-m} = o(1),$$

if

$$p \gg \frac{r^{4r}}{n^{1/(r-1)} \cdot k}. \quad (12)$$

We need to choose p that relations (10), (11) and (12) are all satisfied:

$$p := \frac{r^{4r} \cdot n^{1/2r^2}}{k \cdot n^{1/(r-1)}}.$$

Now we can put together the proof. We started with a p -random subset of the grid $[k]^n$, we remove a point from each of the axis-parallel lines containing at least $r + 1$ points. Also, from each t -tuple of axis-parallel lines intersecting in a (non p -random grid) point, where each line was containing r p -random points we remove a point. The choice of p guarantees that with high probability most of the points were not deleted. After this cleaning, we project this point set into the plane, the way that we do not create new collinear r -tuples. Call the obtained point set S , and write $s := |S|$. We aim to prove that there is no weak ε^* -net T of S of size smaller than $t/5s$. The container method gives that every subset of S of size $3s/4$ contains a collinear r -tuple, i.e., S does not contain a subset of size $s/4$ covering all collinear r -tuples. Note that here we increased $2/3$ to $3/4$, in order to have that this property holds with high probability, and also that the actual size of S might slightly differ from its expectation.

Let $\varepsilon^* = r/s$. Assume that T is a weak ε^* -net of S . Using the switching idea of Alon [2], we find a small ε^* -net, for whose size we have a lower bound. We partition T into three sets, i.e., $T = T_g \cup T_w \cup T_s$, where $T_w = (T - S) \cap v(\mathcal{F})$, $T_s = T \cap S$ and $T_g := T - T_w - T_s$.

As T_g contains only non-grid points, we have that each point of T_g covers at most two lines, hence T_g could be replaced with a set $T'_g \subset S$ covering the same lines that T_g covered, choosing one point from each line, with $2|T_g| \geq |T'_g|$. Each point of T_w could be replaced with at most t points from S to cover the same set of lines, so we could form $T'_w \subset S$ with $t|T_w| \geq |T'_w|$. This means that $T'_g \cup T'_w \cup T_s \subset S$ is an ε^* -net, i.e., has size at least $s/5$. This implies that $|T| > s/5t$.

To summarize, we proved the following result about weak nets. We have a set of size $s \approx p \cdot k^n$, where to cover all the lines containing at least r points we need a set of size at least $s/5\sqrt{r}$. For $\varepsilon^0 > 0$ very small constant we choose k that $r = \log^{1/4} k$, and $s \approx k^{k^r} > 1/\varepsilon^0$. For this choice of k and r we run our proof above. We set $\varepsilon^* := r/s$, and we can get a lower bound on the size of T in function of ε^* :

$$|T| > \frac{s}{5t} = \frac{1}{\varepsilon^*} \cdot \frac{r}{5t} = \frac{1}{\varepsilon^* \cdot 5t} > \frac{1}{10\varepsilon^*} \log \log^{1/10} \frac{1}{\varepsilon^*}.$$

7.1 Decomposing coverage of the plane

Proof of Theorem 2.6: We fix an arbitrary small constant $\gamma > 0$ and $r > r(\gamma)$ (though the method does not require r to be large, the computations are somewhat less tedious under this assumption.) Let $n = 2^{r^4}$. We shall work with the r -uniform hypergraph $\mathcal{F} = \mathcal{F}(n, n, r)$, i.e., $V(\mathcal{F}) = [n]^n$, and $E(\mathcal{F})$ consists of collinear r -tuples of points, where collinearity means that all but one coordinate of the points are the same. We need to compute some parameters of \mathcal{F} :

$$V(\mathcal{F}) = n^n, \quad e(\mathcal{F}) = n^n \cdot \binom{n}{r}, \quad d = d(\mathcal{F}) = r \cdot \binom{n}{r}, \quad \Delta_i = \binom{n-i}{r-i} \quad \text{when } 2 \leq i \leq r.$$

Then we set

$$\varepsilon = 0.5\gamma^r, \quad \tau = \frac{r \cdot 2^r}{n^{1+1/(r-1)}}, \quad p = \frac{r^{4r}}{n^{1+1/(r-1)}}, \quad m = 10\gamma p n^n.$$

Now we apply Theorem 3.1 to \mathcal{F} , where first we have to check that $\Delta(\mathcal{F}, \tau) \leq \varepsilon/(12 \cdot r!)$, we omit the details of the straightforward but tedious computations. We shall obtain a family of containers with at most

$$2^{v(\mathcal{F}) \cdot r^{3r} \cdot n^{-r/(r-1)}}$$

sets, each is spanning at most $\varepsilon \cdot e(\mathcal{F})$ hyperedges, which by Lemma 7.1 means that each set has size at most $\gamma \cdot v(\mathcal{F})$.

Now we sparsen $v(\mathcal{F})$: we keep each vertex with probability p . Now we clean the obtained hypergraph, first we remove vertices from \mathcal{F} from lines with at least $r + 1$ points. Because the expected number of such lines is much less than the expected number of points (because $p \ll n^{-(r+1)/r}$), we remove only few vertices. Additionally, we remove vertices from the remaining random point set, which are on too many, say $r^{5r^2-1} = T$ lines. An application of the Chernoff bound gives that there are not many points removed. We need that in the random hypergraph there are no independent sets of size m , here we have to prove that the expected number of them, in the container sets, is $o(1)$:

$$2^{v(\mathcal{F}) \cdot r^{3r} \cdot n^{-r/(r-1)}} \cdot \binom{\gamma \cdot v(\mathcal{F})}{m} \cdot p^m \leq 2^{v(\mathcal{F}) \cdot r^{3r} \cdot n^{-r/(r-1)}} \cdot \left(\frac{\gamma \cdot v(\mathcal{F}) \cdot p}{10\gamma p \cdot v(\mathcal{F})} \right)^{10\gamma p v(\mathcal{F})} = o(1).$$

Again, the key observation is that the complement of a point set covering all the lines is an independent set, i.e., small independence number implies that the minimum cover should consist of most of the vertices.

Now we can do the standard projection into the plane, and will obtain the required point system.

8 Acknowledgements

The authors are thankful to János Pach who suggested that our method might be applicable in the ε -nets problem. We also thank Noga Alon and Wojciech Samotij for fruitful discussions on the ε -nets problem. Also, we thank for Hong Liu, Adam Zsolt Wagner and the referee for careful reading of the manuscript.

References

- [1] E. Ackerman, J. Pach, R. Pinchasi, R. Radoicic, G. Tóth, A Note on Coloring Line Arrangements, *The Electronic Journal of Combinatorics*, Volume 21, Issue 2 (2014), paper 2.23, 4 pages.
- [2] N. Alon, A non-linear lower bound for planar epsilon-nets, Proc. of the 51th IEEE FOCS (2010), 341–346. Also: *Discrete and Computational Geometry*, **47** (2012), 235–244.
- [3] B. Aronov, E. Ezra, and M. Sharir, Small-size ε -nets for axis-parallel rectangles and boxes, SIAM J. Comput. **39** (2010), no. 7, 3248–3282.

- [4] J. Balogh, H. Liu, and M. Sharifzadeh, The number of subsets of integers with no k -term arithmetic progression, to appear in *Int. Math. Res. Notices*.
- [5] J. Balogh, H. Liu, M. Sharifzadeh and A. Treglown, Sharp bound on the number of maximal sum-free subsets of integers, submitted.
- [6] J. Balogh, R. Morris and W. Samotij, Independent sets in hypergraphs, *J. Amer. Math. Soc.*, **28** (2015), 669–709.
- [7] J. Balogh, R. Mycroft and A. Treglown, A random version of Sperner’s theorem, *Journal of Combinatorial Theory, Series A*, **128** (2014), 104–110.
- [8] B. Bollobás, Random Graphs, *London: Academic Press*, 1985.
- [9] P. Bose, J. Cardinal, S. Collette, F. Hurtado, S. Langerman, M. Korman, and P. Taslakian, Coloring and guarding arrangements, *Discrete Mathematics and Theoretical Computer Science* 15 (2013), 139–154. Also in: 28th European Workshop on Computational Geometry (EuroCG), March 19–21, 2012, Assisi, Perugia, Italy, 89–92.
- [10] M. Collares and R. Morris, Maximum-size antichains in random set-systems, *Random Structures and Algorithms*, to appear.
- [11] P. Erdős: On some metric and combinatorial geometric problems, *Discrete Mathematics*, **60** (1986), 147–153.
- [12] P. Erdős: Problems and results on extremal problems in number theory, geometry, and combinatorics, Proceedings of the 7th Fischland Colloquium, I (Wustrow, 1988), Rostock. Math. Kolloq., No. 38 (1989), 6–14.
- [13] P. Erdős: Some old and new problems in combinatorial geometry, Applications of discrete mathematics (Clemson, SC, 1986), 32–37, SIAM, Philadelphia, PA, 1988.
- [14] Z. Füredi, Maximal independent subsets in Steiner systems and in planar sets, *SIAM J. Discrete Math.*, 4 (1991), 196–199.
- [15] Z. Füredi and J. Pach, Traces of finite sets: extremal problems and geometric applications, in: Extremal problems for finite sets (Visegrád, 1991), 251–282, Bolyai Soc. Math. Stud., 3, János Bolyai Math. Soc., Budapest, 1994.
- [16] H. Furstenberg and Y. Katznelson, A density version of the Hales-Jewett theorem for $k = 3$, Graph Theory and Combinatorics (Cambridge, 1988), *Discrete Math.*, **75** (1989), 227–241.
- [17] H. Furstenberg and Y. Katznelson, A density version of the Hales-Jewett theorem, *J. Anal. Math.*, **57** (1991), 64–119.
- [18] D. Haussler and E. Welzl, ε -nets and simplex range queries, *Discrete and Computational Geometry*, (1987), 127–151.

- [19] J. Kollár, Szemerédi-Trotter-type theorems in dimension 3, *Advances in Mathematics*, **271** (2015), 30–61.
- [20] J. Komlós, J. Pach, and G. Woeginger, Almost tight bounds for ε -nets, *Discrete Comput. Geom.*, **7** (1992), 163–173.
- [21] P. Mani-Levitska and J. Pach, J. Decomposition problems for multiple coverings with unit balls, (1988), manuscript.
- [22] J. Matousek, Lectures in Discrete Geometry, Graduate Texts in Mathematics, 212. Springer-Verlag, New York, 2002. xvi+481 pp. ISBN: 0-387-95373-6
- [23] J. Matousek, R. Seidel and E. Welzl, How to net a lot with little: Small ε -nets for disks and halfspaces, In Proc. 6th Annu. ACM Sympos. Comput. Geom., 16–22, 1990.
- [24] R. Nenadov and A. Steger, A short proof of the Random Ramsey theorem, *Combinatorics, Probability, and Computing*, **25** (2016), 130–144.
- [25] J. Pach and P. Agarwal, Combinatorial Geometry, John Wiley & Sons, Inc. (1995) DOI: 10.1002/9781118033203.
- [26] J. Pach and G. Tardos, Tight lower bounds for the size of epsilon-nets *J. Amer. Math. Soc.* **26** (2013), 645–658.
- [27] J. Pach, G. Tardos and G. Tóth, Indecomposable Coverings, *Canad. Math. Bull.*, **52** (3) (2009), 451–463.
- [28] K. T. Phelps and V. Rödl, Steiner triple systems with minimum independence number, *Ars Combin.*, **21** (1986), 167–172.
- [29] D.H.J. Polymath, A new proof of the density Hales-Jewett theorem. *Ann. of Math.*, **175** (2012), no. 3, 1283–1327.
- [30] D.H.J. Polymath, Density Hales-Jewett and Moser Numbers, in: An Irregular Mind (eds. Bárány et al), Bolyai Society Mathematical Studies V 21, Springer, 2010, 689–753.
- [31] V. Rödl, A. Ruciński and M. Schacht, An exponential-type upper bound for Folkman numbers, to appear in *Combinatorica*.
- [32] D. Saxton and A. Thomason, Hypergraph containers, *Inventiones mathematicae*, (2015), 1–68.